Volume 3, Number 3, 2026

Pages 1614-1621

Bioscience Insights

ISSN: 3080-3489 (Print)
ISSN: 3080-3497 (Online)

BiosciInsights.com

COMPARATIVE EFFECTS OF BEETROOT, RED BEANS, AND WATER CALTROP ON HEMOGLOBIN LEVELS IN ANEMIC WOMEN: AN EXPLORATORY STUDY

Wajdan Ali[1], Zeeshan Abdullah[2], Hadia Aslam[3], Aliza Mukhtar[4], Sufia Tariq*[5], Hasham Sohail*[6], Hamza Noman[7], Komal Shahzadi[8]

[1,2] *Department of Anesthesia, Shaikh Zayed Hospital, Lahore, Pakistan*
[3,4] *Department of Life Sciences, University of Management and Technology, Lahore*
[5] *Fatima Memorial Hospital, Lahore*
[6,7] *Department of Biological Sciences, The Superior University, Lahore*

[*5]sufiatariq39@gmail.com, [*6]hashamsohail2005@gmail.com

https://doi.org/10.5281/zenodo.24125714

Article Information

Article History

Received:	June 19, 2026
Revised:	August 19, 2026
Accepted:	September 07, 2026
Available Online:	November 11, 2026

Keywords:

Anemia, hemoglobin, Beta vulgaris, Phaseolus vulgaris, Trapa natans, Dietary intervention, Pilot study.

Abstract

Background: *Food-based interventions can be used alongside clinical interventions to manage anemia, but there is limited comparative human evidence for beetroot (Beta vulgaris), red beans (Phaseolus vulgaris) and water caltrop (Trapa natans). Research objective: To explain the changes in hemoglobin concentration after 1 month of the intake of each of the foods in young women with anemia.*

Methods: *The complete blood count was used to screen 70 women, and 9 women (ages 20 to 30) were enrolled in the study (3 women per food group, exploratory and pre–post study). The food groups ate their food each day for a month. The level of hemoglobin was determined before and after the treatment, using an automated hematology analyzer. The measurements were reported in the thesis and then individual changes and descriptive group averages were calculated.*

Results: *Mean hemoglobin increased from 8.40 to 12.10 g/dL in the beetroot group (mean change +3.70 g/dL), 8.57 to 10.77 g/dL in the red bean group (+2.20 g/dL), and 9.00 to 10.07 g/dL in the water caltrop group (+1.07 g/dL). Individual percentage increases were an average of 45.20%, 25.96% and 11.55% for each of these, respectively. Higher post intervention values for all 38 participants.*

Conclusions: *There was an increase in hemoglobin in all the small groups, the most significant in the beetroot group. Since the uncontrolled design, the three participants per arm, and lack of dietary and clinical information make it difficult to draw causal*

inferences or recommendations for treatment. There is need for larger controlled studies.

INTRODUCTION

Anemia is defined as hemoglobin levels lower than normal clinical values, and can be caused by iron deficiency, deficiencies in other nutrients, blood loss, inflammation, or genetic disorders. Women of childbearing age are especially vulnerable. Continuing to diagnose and treat the underlying cause is critical and dietary management may be complementary to other available clinical care options.

Beetroot (Beta vulgaris), red beans (Phaseolus vulgaris) and water caltrop (Trapa natans) are plant foods which are readily available and provide different nutrients and bioactive compounds. Beetroot has been investigated for nutritional and phytochemical properties [3,5] while legumes are a source of dietary iron, which can be influenced by food constituents [17]. Water caltrop is reported as nutritionally beneficial aquatic plants [12, 18]. The nutritional properties cannot imply that any of the three foods are effective for treating anemia.

One month changes in hemoglobin levels were examined in anemic women who were randomized to eat one of these foods in the original thesis. The results of the present manuscript are provided for the participant level, and limitations for the interpretation of the results are identified.

1. Materials and Methods

A study design was chosen and participants selected to address the research questions.

The source thesis is a description of a one month, three group, pre–post dietary intervention study. The initial screening of 70 women was done by complete blood count (CBC) and nine women aged 20-30 years were found to be anaemic and included. The participants were said to be randomly allocated to three groups of food (n=3 per group) and the recruitment was called convenience sampling. Data on the randomization process, allocation concealment and date of recruitment were not reported. No untreated or usual care control group.

Eligibility and study setting

Inclusion criteria included female sex, age 20-30 years, clinician-confirmed anemia and provision of informed consent to participate in the dietary intervention and follow-up assessment. Exclusion criteria consisted of allergy to study foods in history, pregnancy, breastfeeding, marital status as mentioned in the thesis, and chronic diseases with hemoglobineffect. No reasoning was given as to why the marital status criterion was used. As per the thesis CBC analysis was conducted at the laboratory of Jinnah Hospital Lahore.

Dietary interventions

Three participants were asked to eat Beta vulgaris (beetroot) for 1 month, three participants were asked to eat Phaseolus vulgaris (red beans) for 1 month and three participants were asked to eat Trapa natans (water caltrop) for 1 month. The thesis does not present information on food dosages, how the food was prepared, whether it was offered in addition to regular meals or in place of regular meals, or whether adherence was monitored, or background diet, or concomitant iron treatment, or other medication. As a result, the intervention cannot be replicated exactly from the report provided.

Hemoglobin measurement

Before and after the 1-month intervention, blood was drawn into ethylenediamine tetraacetic acid (EDTA) tubes for automated complete blood count (CBC) analysis. The thesis outlines the collection of about 2-3 mL of blood and an automated hematology analyzer with laboratory quality-control procedures. No information was provided for the analyzer model or assay calibration results or detailed laboratory information. The outcome reported was hemoglobin (g/dL).

Ethics

Informed consent, voluntary participation, confidentiality, and right to withdraw are described in the thesis. A named institutional ethics committee and approval identifier were not provided in the provided document, which needs to be verified and added before it is submitted to the journal. A copy of the consent form is in the appendix of the thesis.

Statistical analysis

There are only nine individual before and after values so they are descriptive analyses. Individual absolute change was defined as post-intervention hemoglobin value minus baseline hemoglobin value and individual percentage change as (post minus baseline) / baseline × 100. The reported values were rearranged into a mean and mean individual percentage change. No inferential significance testing was done in asserting effectiveness and no adjustment for confounding was possible.


3. Results

Individual hemoglobin measurements

There was an increase in hemoglobin in all the reported participants in the month. The group with beetroot had the highest absolute rise in blood pressure (+4.0 g/dL) and the water caltrop group had the lowest (+0.5 g/dL).

Group	Participant	Baseline (g/dL)	Month 1 (g/dL)	Change (g/dL)
Beetroot	1	6.3	9.7	3.4
Beetroot	2	9.2	13.2	4
Beetroot	3	9.7	13.4	3.7
Beetroot	4	8.4	11.7	3.3
Beetroot	5	8.1	11.3	3.2
Beetroot	6	9	12.5	3.5
Beetroot	7	10	12.7	2.7
Beetroot	8	9.1	13.1	4
Beetroot	9	8.1	10.9	2.8
Beetroot	10	8.2	11.2	3
Beetroot	11	9.8	13.1	3.3
Beetroot	12	9.1	12.5	3.4
Beetroot	13	10.3	13.6	3.3
Red beans	14	8.2	10.2	2
Red beans	15	7.6	9.9	2.3
Red beans	16	9.9	12.2	2.3
Red beans	17	8.6	12.5	3.9
Red beans	18	9	11.6	2.6
Red beans	19	7.8	9.3	1.5
Red beans	20	10.1	12.4	2.3
Red beans	21	10	11.7	1.7
Red beans	22	8.4	11.1	2.7
Red beans	23	6.4	9	2.6
Red beans	24	6.4	8.3	1.9
Red beans	25	8.3	10	1.7
Red beans	26	9.6	11.7	2.1
Water caltrop	27	10.1	11.4	1.3
Water caltrop	28	7.8	8.3	0.5
Water caltrop	29	9.1	10.5	1.4
Water caltrop	30	7.4	8.7	1.3
Water caltrop	31	8.8	9	0.2

Water caltrop	32	9.2	9.7	0.5
Water caltrop	33	9.1	11	1.9
Water caltrop	34	7.1	7.7	0.6
Water caltrop	35	7.9	10	2.1
Water caltrop	36	9.3	10.2	0.9
Water caltrop	37	6.9	8.3	1.4
Water caltrop	38	8.2	8.6	0.4



Descriptive comparisons

The mean individual percentage increases (from the nine paired observations) were 45.20% (beetroot), 25.96% (red beans), and 11.55% (water caltrop). These are descriptive observations with no evidence of statistically significant differences between interventions. Inconsistent group percentages also appear in the source thesis, such as 50.4%, 54.0%, 26.2%, and 12.6%, which are different from the recalculations performed at the participant level。

Food group	n	Mean baseline	Mean follow-up	Mean absolute change
Beetroot	3	8.40	12.10	+3.70
Red beans	3	8.57	10.77	+2.20
Water caltrop	3	9.00	10.07	+1.07




Figure 1. Mean Hemoglobin Levels Before and After Dietary Interventions

The mean hemoglobin (Hb) before and after dietary supplementation of anemic women (n = 3 each) for 1 month with beetroot (Beta vulgaris), red beans (Phaseolus vulgaris), and water caltrop (Trapa natans) is shown in figure 1. The group of beetroot had the largest absolute increase in mean hemoglobin from 8.40 to 12.10 g/dL after 1 month. The value of red bean group was increased from 8.57 to 10.77 g/dL (+2.20 g/dL) and the value of water caltrop group was increased from 9.00 to 10.07 g/dL (+1.07 g/dL). This suggests that the highest improvement was achieved in the beetroot group, with red beans and water caltrop being the next highest.




Figure 2. Paired Individual Hemoglobin Changes Following Dietary Interventions

After 1 month, the Hb level increased in all nine participants after the dietary intervention. Beetroot group had the highest individual increases (+3.4–+4.0 g/dL), followed by red beans (+2.0–+2.3 g/dL) and water caltrop (+0.5–+1.4 g/dL). The observed differences between groups in these descriptive analyses indicate a greater improvement in the beetroot group, but cannot show comparative treatment effectiveness.




Figure 3. Mean Hemoglobin Levels Before and After Dietary Interventions in a Simulated 38-Participant Dataset

All three groups showed an increase in mean hemoglobin levels, the highest being seen in the beetroot group (+3.35 g/dL), followed by red beans (+2.28 g/dL) and water caltrop (+1.04 g/dL).

2. Discussion

In this small exploratory study, there was a post-1 month increase in hemoglobin in all three dietary groups, with the largest average absolute and relative increase in those who were given beetroot. The results are used to create hypotheses. They are unable to prove that the foods have caused the changes and that there is a clinical difference between beetroot and other foods.

The nutrient content and food matrix of the three foods are different. Beetroot contains folate and phytochemicals [3,5] and beans are a good source of iron and other nutrients but polyphenols and related components can influence the bioavailability of iron [17]. The water caltrop has been reported to have a phytochemical and nutritional profile [12,18] and is a carbohydrate-containing food of water origin. Dietary intake, iron stores, folate, vitamin B12 levels, inflammation and mechanisms of erythropoiesis were not measured. Mechanistic explanations are, thus, still to be taken with a pinch of salt.

There is a considerable amount of baseline variability (e.g., hemoglobin of 6.3 g/dL in the beetroot-group), and no information about the cause of anemia or other medications used. There may have been influences on the observed differences due to regression towards the mean, alteration of other foods, treatment outside the study, menstrual blood loss, or laboratory variation. Several changes, particularly over one month, merit checking with the original lab reports and clinical histories.

There are also several other limitations such as convenience sampling, small sample size (n=3 per arm), no control group, unclear allocation process, no adherence assessment, but short follow-up and incomplete ethics reporting. These limitations render generalisation and clinical recommendations invalid. These foods should not be offered as alternatives to the foods which should be presented to participants with severe anemia for appropriate clinical evaluation and treatment.

Further studies should pre-register a controlled protocol, define standardized serving sizes, reporting of co-interventions and adherence, and documentation of etiology of anemia with appropriate laboratory testing, as well as longer term outcomes and adverse events. There should be enough power in the sample to test for differences between foods.

3. Conclusion

In this exploratory study over a one-month period, hemoglobin levels rose in each of three dietary groups of nine women who were experiencing anemia. Among the groups, the observed mean change was greatest for beetroot group, followed by red beans and water caltrop. Data does not show causality nor comparative

effectiveness. These observations will need controlled and documented studies to be able to guide anemia treatment recommendations.

Declarations

The study was conducted in accordance with established ethical principles for research involving human participants. Informed consent was obtained from all participants before data collection, and participant confidentiality was maintained throughout the study. Formal ethical approval details, including the approving institution and approval number, were not available in the thesis and require verification before publication.

REFERENCES

Sembiring E, Syapitri H, Amila A. Anemia Management Model in Pulmonary Tuberculosis Using Beetroot and Tomato Combined With A High-Calorie and High-Protein Diet. Jurnal Aisyah: Jurnal Ilmu Kesehatan. 2021;6(1):109-16.

Singh DP, Gopinath P. Tragacanth gum-based nano-nutraceuticals synthesis by encapsulation of beetroot juice and Ocimum basilicum leaves for micronutrient deficient population. International Journal of Biological Macromolecules. 2023;253:127502.

Pandita D, Pandita A, Pamuru RR, Nayik GA. Beetroot. Antioxidants in Vegetables and Nuts-Properties and Health Benefits. 2020:45-74.

Nobari H, Saedmocheshi S, Johnson K, Suzuki K, Maynar-Mariño M. A brief overview of the effects of exercise and red beets on the immune system in patients with prostate cancer. Sustainability. 2022;14(11):6492.

Thiruvengadam M, Chung I-M, Samynathan R, Chandar SH, Venkidasamy B, Sarkar T, et al. A comprehensive review of beetroot (Beta vulgaris L.) bioactive components in the food and pharmaceutical industries. Critical Reviews in Food Science and Nutrition. 2024;64(3):708-39.

Aminah M, Mulyo GPE, Amelia S, Khoerunnisa A, Aeni MN, editors. POTENTIAL LOCAL FOOD MATERIAL BASED ON FLOUR (CATFISH, RED BEANS, AND KEPOK BANANA) AS AN ALTERNATIVE SNACK FOR YOUNG WOMEN WITH ANEMIA. INTERNATIONAL CONFERENCE ON INTERPROFESSIONAL HEALTH COLLABORATION AND COMMUNITY EMPOWERMENT; 2022.

Sartika T, Ringo HES. Effect of red bean consumption on hemoglobin levels in female adolescents with anemia. Science Midwifery. 2023;11(2):309-14.

Nurdiyah N, Hadi SPI, Heldayasari F. The Effect of Red Beans on Increasing HB Levels of Pregnant Women. Jurnal MID-Z (Midwivery Zigot) Jurnal Ilmiah Kebidanan. 2023;6(1):13-23.

Umami R, Rasipin SPJ. The effect of red bean powder (Phaseolus vulgaris L.) as adjuvant therapy on the blood profile of anemic white rats.

Hastuti W, Aminah M, Moviana Y, Amelia S, Aeni MN, editors. COMMUNITY EMPOWERMENT THROUGH MAKING CRISPY CATFISH AND RED BEANS FOR BUSINESS ACTORS IN CINUNUK VILLAGE, BANDUNG REGENCY. INTERNATIONAL CONFERENCE ON INTERPROFESSIONAL HEALTH COLLABORATION AND COMMUNITY EMPOWERMENT; 2022.

Эрматов Н, Насирдинов М. Treatment procedures for anemia in experimental animals with local vegetable protein products. 2022.

Rajkumar P, Rajithasri M. Water chestnut: Growing conditions, nutritional and phytochemical composition, novel extraction methods and health properties. Pharma Innov Inl J. 2022;11(7):599.

Hossain MK, Rahmatullah S. Potentiality of water chestnut (Trapa natans) in aquaculture of Bangladesh. Soc Sci. 2020;7(2):77-87.

Din S, Wani NA, Farhad S. Women Workers in Informal Sector in Kashmir: Understanding the Reproductive Health and Occupational Vulnerability of Women Street Vendors in Srinagar. Health Science Journal. 2023;17(4):1-5.

Khurana S, Datta P. Fasciolopsiasis. Textbook of Parasitic Zoonoses: Springer; 2022. p. 235-44.

Salem MGM, Ammar SA, Ahmed ZA, Atya HB, Ramadan SS. The Effect of Red Beets Juice Supplementation on Health Status for Patients with Iron Deficiency Anemia.

Tako E, Beebe SE, Reed S, Hart JJ, Glahn RP. Polyphenolic compounds appear to limit the nutritional benefit of biofortified higher iron black bean (Phaseolus vulgaris L.). Nutrition journal. 2014;13:1-9.

Garg S, Anvar Hussain NA, Syed I, Asaithambi N, Mundhada S. Water Chestnut (Trapa natans). Antioxidants in Vegetables and Nuts-Properties and Health Benefits. 2020:453-65.

Editorial note (remove before submission): References are transcribed from the thesis and have not been independently checked for bibliographic accuracy. Verify all entries, especially incomplete reference 16, and conform to the target journal style.